

Volume 19, Issue 1, April 2024

p-ISSN : 1829-7005

e-ISSN : 2540-8836

The Indonesian JOURNAL *of* PUBLIC HEALTH

Scientific Journal of Public Health



The Indon. J of PH	Vol. 19	Issue. 1	Page 1-195	Surabaya April 2024	p-ISSN : 1829-7005 e-ISSN : 2540-8836
-----------------------	---------	----------	---------------	------------------------	--

Vol. 21 No. 1 (2026): THE INDONESIAN JOURNAL OF PUBLIC HEALTH



Published: 02-04-2026

Editorial

MAPPING VARIOUS HEALTH CONCEPTS IS CRUCIAL IN UNDERSTANDING HEALTH PROBLEMS TOWARDS BETTER ACADEMIC AND PRACTICAL PERFORMANCE

 Kuntoro Kuntoro ⁽¹⁾

 1-

(1) Department of Epidemiology, Biostatistics, Population Studies and Health Promotion, Public Health Faculty, Universitas Airlangga, Indonesia

2

 Abstract : 160

 XML : 0 PDF : 343

 XML

 PDF

 DOI : 10.20473/ijph.v21i1.2026.1-2

Articles



KNOWLEDGE, ATTITUDE, AND PRACTICE TOWARD PREVENTION AND MANAGEMENT OF MEDICATION ERRORS AMONG PATIENTS: A CROSS-SECTIONAL STUDY

(1) Department of Pharmacy, Faculty of Health Sciences, Jenderal Soedirman University, Purwokerto, Indonesia, Indonesia ,

(2) Department of Pharmacy, Faculty of Health Sciences, Jenderal Soedirman University, Purwokerto, Indonesia, Indonesia ,

(3) Department of Pharmacy, Faculty of Health Sciences, Jenderal Soedirman University, Purwokerto, Indonesia, Indonesia ,

(4) Department of Pharmacy, Faculty of Health Sciences, Jenderal Soedirman University, Purwokerto, Indonesia, Indonesia


 Abstract : 422

 XML : 23 PDF : 654

 XML

 PDF

 DOI : 10.20473/ijph.v21i1.2026.3-17



IS SLEEP DURATION CORRELATE TO THE ANTHROPOMETRIC MEASUREMENTS AND INSULIN LEVELS IN OBESE ADOLESCENTS? INVESTIGATION WITH A CROSS-SECTIONAL STUDY

 Eva Ardianah ⁽¹⁾, Nur Aisiyah Widjaja ⁽²⁾, Tausyiah Rohmah Noviyanti ⁽³⁾, Yoppi Yeremia Alexander ⁽⁴⁾, Yasmine Nurfirdaus ⁽⁵⁾, Rino Tryanto Keya ⁽⁶⁾, Edi Hermanto ⁽⁷⁾, Iitdrie ⁽⁸⁾, Welldelin Yufuria Christiansi ⁽⁹⁾



18-

29

(1) Master of Public Health, Universitas Airlangga, Surabaya, Indonesia, Indonesia ,

(2) Department of Child Health, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia, Indonesia ,

(3) Master of Public Health, Universitas Airlangga, Surabaya, Indonesia, Indonesia ,

(4) Master of Public Health, Universitas Airlangga, Surabaya, Indonesia, Indonesia ,

(5) Master of Public Health, Universitas Airlangga, Surabaya, Indonesia, Indonesia ,

(6) Master of Public Health, Universitas Airlangga, Surabaya, Indonesia, Indonesia ,

(7) Master of Public Health, Universitas Airlangga, Surabaya, Indonesia, Indonesia ,

(8) Master of Public Health, Universitas Airlangga, Surabaya, Indonesia, Indonesia ,

(9) Master of Public Health, Universitas Airlangga, Surabaya, Indonesia, Indonesia

 Abstract : 308

 PDF : 399 XML : 23

 PDF

 XML

 DOI : 10.20473/ijph.v21i1.2026.18-29



THE IMPACT OF COVID-19 ON ELDERLY DAY CARE A CASE STUDY IN BANGKOK, THAILAND

Nakarait Na Pattalung ⁽¹⁾, Maniga Thongkong ⁽²⁾

30-42

(1) Faculty of Management Science, Suan Dusit University, Bangkok, 10300 Thailand, Thailand ,

(2) ARC Condo, Vibhavadi Road, Bang Khen, Lak Si, Bangkok, 10210 Thailand, Thailand

Abstract : 183

XML : 17 PDF : 232



DOI : 10.20473/ijph.v21i1.2026.30-42



COGNITIVE UNDERPINNING OF POST TRAUMATIC STRESS DISORDER AMONG AFGHAN AND KYRGYZ STUDENTS: AN EMPIRICAL INVESTIGATION

Azizuddin Khan ⁽¹⁾, Elena Molchanova ⁽²⁾, Zarlisht Sarmast ⁽³⁾, Farangis Dehnavi ⁽⁴⁾, Eshquvvatov To'liqinjon Eshquvvatovich ⁽⁵⁾, Umrah Beg ⁽⁶⁾

43-56

(1) Psychophysiology Laboratory, Department of Humanities and Social Sciences, Indian Institute of Technology Bombay, Mumbai-400076, Maharashtra, India, India ,

(2) Division of Social Sciences, Department of Psychology, American University of Central Asia (office 315), 7/6 Aaly Tokombaev Street, Bishkek, Kyrgyz Republic, Kyrgyzstan ,

(3) Global Fellows Program Coordinator, SUN, AUCA, Bard – CCE, Kyrgyzstan ,

(4) Psychophysiology Laboratory, Department of Humanities and Social Sciences, Indian Institute of Technology Bombay, Mumbai-400076, Maharashtra, India, India ,

(5) Head of the Department of “Pre-school and primary education, sports”, Kattakorgan Branch of Samarkand State University, Uzbekistan, Uzbekistan ,

(6) Psychophysiology Laboratory, Department of Humanities and Social Sciences, Indian Institute of Technology Bombay, Mumbai-400076, Maharashtra, India, India

Abstract : 215

XML : 15 PDF :



THE IMPACT OF EARMUFF USE ON HEARING FUNCTION IMPAIRMENT AND FATIGUE: A TIME SERIES STUDY AMONG WEAVING WORKERS IN SURAKARTA, INDONESIA

Haris Setyawan ⁽¹⁾, Nisa Nur Kusuma ⁽²⁾, Ratih Andhika Akbar Rahma ⁽³⁾, Tofan Agung Eka Prasetya ⁽⁴⁾

57-71

(1) Occupational Health and Safety Study Program, Vocational School, Universitas Sebelas Maret Surakarta, Indonesia, Indonesia ,


(2) Health Information Management Study Program, Politeknik Assalaam Surakarta, Indonesia, Indonesia ,

(3) Occupational Health and Safety Study Program, Universitas Darussalam Gontor, Indonesia, Indonesia ,

(4) Health Department, Faculty of Vocational Studies, Universitas Airlangga, Campus B, Surabaya , Indonesia

Abstract : 207

PDF : 289 XML : 22



MALNUTRITION RISK FACTORS BASED ON SLEEP DURATION AND PHYSICAL ACTIVITY: STUDIES IN BANGLADESH AND INDONESIA

Catur Wulandari ⁽¹⁾, Anugrah Linda Mutiarani ⁽²⁾, Pratiwi Hariyani Putri ⁽³⁾, Sa'bania Hari Raharjeng ⁽⁴⁾

72-84

(1) Departement of Nutrition, Faculty of Health, Universitas Nahdlatul Ulama Surabaya, Surabaya, Indonesia, Indonesia ,

(2) Departement of Nutrition, Faculty of Health, Universitas Nahdlatul Ulama Surabaya, Surabaya, Indonesia, Indonesia ,

(3) Departement of Nutrition, Faculty of Health, Universitas Nahdlatul Ulama Surabaya, Surabaya, Indonesia, Indonesia ,



NAVIGATING CROSS-BORDER HEALTHCARE POLICIES AT THE SARAWAK-WEST KALIMANTAN BORDER

 Eva Novianty Rachmat ⁽¹⁾, Dolly Paul Carlo ⁽²⁾, Helmy Hazmi ⁽³⁾

 85-100

(1) Human Resource Development Agency, Ministry of Transportation, Republic of Indonesia, Indonesia ,

(2) Faculty of Social Sciences and Humanities, Universiti Malaysia Sarawak, Malaysia , Malaysia ,

(3) Faculty of Medicine and Health Sciences, Universiti Malaysia Sarawak, Malaysia, Malaysia



THE RISK FACTORS AND BEHAVIORS ASSOCIATED WITH HOUSE DUST MITE-RELATED ALLERGIC RESPIRATORY DISEASES: A HOSPITAL-BASED STUDY

 Natalia Erica Jahja ⁽¹⁾, Zahrah Hikmah ⁽²⁾

(1) Allergy and Immunology Fellowship Program, Department of Child Health, Faculty of Medicine, Universitas Airlangga, Surabaya 60132, Indonesia; Dr. Soetomo General Academic Hospital, Surabaya 60285, Indonesia; Department of Pediatrics, Petrokimia Gresik Hospital, Gresik 61119, Indonesia, Indonesia ,


 Abstract : 150

 XML : 15 PDF : 227

 XML

 PDF

 DOI : 10.20473/ijph.v21i1.2026.101-112



CONCEPT MAPPING CARE PLAN (CMCP): AN APPROACH TO IMPROVING THE ACADEMIC PERFORMANCE AMONG NURSING STUDENTS IN THE CLINICAL SETTING

 Rusnani Ab Latif ⁽¹⁾, Wan Ismahanisa Ismail ⁽²⁾, Yoyok Bkti Prasetyo ⁽³⁾, Abd Nasir ⁽⁴⁾, Jamilah Shaari ⁽⁵⁾

 113-125

(1) Faculty of Health Sciences, Universiti Teknologi MARA, Cawangan Pulau Pinang, Campus Bertam,13200, Kepala Batas, Pulau Pinang, Malaysia, Malaysia ,

(2) Faculty of Health Sciences, Universiti Teknologi MARA, Cawangan Pulau Pinang, Campus Bertam,13200, Kepala Batas, Pulau Pinang, Malaysia, Malaysia ,

(3) Faculty of Health Sciences, University of Muhammadiyah Malang, Indonesia, Indonesia ,

(4) Faculty of Vocational Studies, Universitas Airlangga, Surabaya, Indonesia, Indonesia ,

(5) School of Health Sciences, Health Campus Universiti Sains Malaysia, 16150 Kubang Kerian, Kelantan, Malaysia, Malaysia

 Abstract : 129

 PDF : 203 XML : 15

 PDF

 XML

 DOI : 10.20473/ijph.v21i1.2026.113-125



IMPACT OF ATHLETIC PARTICIPATION ON THE BODY COMPOSITION OF COLLEGE STUDENTS


- (1) Department of Physical Education, Faculty of Humanities and Social Sciences, Nims University Rajasthan, 139 Jaipur, India, India ,
(2) Department of Physical Education, Faculty of Humanities and Social Sciences, Nims University Rajasthan, Jaipur, India, India ,
(3) Department of Education in Social Science, NCERT, New Delhi, India, India ,
(4) Department of Physical Education and Yoga, ICAI University, Tripura, India, India

 Abstract : 129

 PDF : 198 XML : 24



 DOI : 10.20473/ijph.v21i1.2026.126-139



BIBLIOMETRIC ANALYSIS OF AIR POLLUTION'S IMPACT ON CHILDREN WITH ASTHMA: INSIGHTS FROM A DECADE OF RESEARCH (2012-2022)

 Ichtiarini Nurullita Santri ⁽¹⁾, Lalu Muhammad Irham ⁽²⁾, Musfirah Musfirah ⁽³⁾, Yuniar Wardani ⁽⁴⁾, Khoiriyah Isni ⁽⁵⁾, Yohane Vincent Abero Phiri ⁽⁶⁾, Ahmad Faizal Rangkuti ⁽⁷⁾, Nopryan Ekadinata ⁽⁸⁾, Wirawan Adikusuma ⁽⁹⁾ 

140-

- (1) Faculty of Public Health, Universitas of Ahmad Dahlan, Yogyakarta, Indonesia, Indonesia ,
(2) Faculty of Pharmacy, Universitas of Ahmad Dahlan, Yogyakarta, Indonesia, Indonesia ,
(3) Faculty of Public Health, Universitas of Ahmad Dahlan, Yogyakarta, Indonesia, Indonesia ,
(4) Faculty of Public Health, Universitas of Ahmad Dahlan, Yogyakarta, Indonesia, Indonesia ,
(5) Faculty of Public Health, Universitas of Ahmad Dahlan, Yogyakarta, Indonesia, Indonesia ,
(6) University at Buffalo, Department of Epidemiology and Environmental Health (EEH), Buffalo NY, USA, United States ,
(7) Faculty of Public Health, Universitas of Ahmad Dahlan, Yogyakarta, Indonesia, Indonesia ,
(8) School of Nursing, Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Yogyakarta, Yogyakarta, Indonesia, Indonesia ,
(9) Department of Pharmacy, Universitas Muhammadiyah Mataram, Mataram, Indonesia; Research Center for Computing, National Research and Innovation Agency (BRIN), Cibinong Science Center, Cibinong, Indonesia, Indonesia


152

 Abstract : 135

 PDF : 294 XML : 19



 DOI : 10.20473/ijph.v21i1.2026.140-152



EARLY DETECTION OF HYPERTENSION RISK: A SURVIVAL ANALYSIS OF BLOOD PRESSURE IN AGING POPULATIONS IN SEMARANG

Putri Tiara Rosha ⁽¹⁾, Ayu Rahadiyanti ⁽²⁾, Dina Nur Anggraini Ningrum ⁽³⁾, Novia Fariqothul Qudsyiah ⁽⁴⁾, Firly Azra Ghassanie ⁽⁵⁾, Emilia Indri Artanti ⁽⁶⁾, Irna Mufidatul Himmah ⁽⁷⁾, Florensia Indah Setyowati ⁽⁸⁾, Yuniar Dwi Prastika ⁽⁹⁾

153-167

(1) Public Health Department, Faculty of Medicine, Universitas Negeri Semarang, Indonesia , Indonesia ,

(2) Nutrition Department, Faculty of Medicine, Universitas Diponegoro, Indonesia, Indonesia ,

(3) Public Health Departement, Faculty of Medicine, Universitas Negeri Semarang, Indonesia , Indonesia ,

(4) Public Health Departement, Faculty of Medicine, Universitas Negeri Semarang, Indonesia, Indonesia ,

(5) Public Health Departement, Faculty of Medicine, Universitas Negeri Semarang, Indonesia , Indonesia ,

(6) Public Health Departement, Faculty of Medicine, Universitas Negeri Semarang, Indonesia, Indonesia ,

(7) Public Health Departement, Faculty of Medicine, Universitas Negeri Semarang, Indonesia , Indonesia ,

(8) Medicine Department, Faculty of Medicine, Universitas Negeri Semarang, Indonesia, Indonesia ,

(9) Public Health Departement, Faculty of Medicine, Universitas Negeri Semarang, Indonesia , Indonesia


Abstract : 144

XML : 17 PDF : 222

XML

PDF

DOI : 10.20473/ijph.v21i1.2026.153-167



DETERMINANTS OF TIME TO PULMONARY TUBERCULOSIS DEVELOPMENT IN TYPE II DIABETES MELLITUS PATIENTS AT A COMMUNITY HOSPITAL

Nichanan Nakwihok ⁽¹⁾, Sunisa Ruangmuan ⁽²⁾, Sutee Nuket ⁽³⁾, Somkiattiyos Woradet ⁽⁴⁾, Bhunyabhadh Chaimay ⁽⁵⁾, Kawintida Jeenuang ⁽⁶⁾

167-182

(1) Faculty of Health and Sports Science, Thaksin University, Phatthalung campus, Phatthalung Province, 93210 Thailand, Thailand ,

(2) Faculty of Health and Sports Science, Thaksin University, Phatthalung campus, Phatthalung Province, 93210 Thailand, Thailand ,


- (3) Pak Phanang Hospital, Nakhon Si Thammarat Province, Thailand, Thailand ,
 (4) Faculty of Health and Sports Science, Thaksin University, Phatthalung Campus, Phatthalung Province, Thailand, Thailand ,
 (5) Faculty of Health and Sports Science, Thaksin University, Phatthalung Campus, Phatthalung Province, Thailand, Thailand ,
 (6) Faculty of Health and Sports Science, Thaksin University, Phatthalung Campus, Phatthalung Province, Thailand, Thailand

 Abstract : 115

 XML : 15 PDF : 230



 DOI : 10.20473/ijph.v21i1.2026.168-182



PORNOGRAPHY EXPOSURE AND SEXUAL BEHAVIOR AMONG COLLEGE STUDENTS AT UNIVERSITAS JEMBER

 N. Dyandra Agnefa Argadini ⁽¹⁾, Ni mal Baroya ⁽²⁾, Devi Arine Kusumawardani ⁽³⁾

 183-194

(1) Undergraduate Public Health Study Program, Faculty of Public Health, Universitas Jember, Indonesia, Indonesia ,

(2) Department of Epidemiology, Biostatistics and Population Studies, Indonesia ,


(3) Faculty of Public Health, Universitas Jember, Indonesia

 Abstract : 94

 XML : 48 PDF : 192



 DOI : 10.20473/ijph.v21i1.2026.183-194



REPRODUCTIVE HEALTH KNOWLEDGE AMONG ADOLESCENTS WITH INTELLECTUAL DISABILITY EDUCATIONAL AND INTERVENTION CHALLENGES

(1) Department of Public Health, Universitas Negeri Semarang, Semarang, Indonesia; Poltekkes Kemenkes Semarang, Semarang, Indonesia, Indonesia ,

(2) Department of Public Health, Universitas Negeri Semarang, Semarang, Indonesia, Indonesia ,

(3) Department of Public Health, Universitas Negeri Semarang, Semarang, Indonesia, Indonesia ,

(4) Department of Public Health, Universitas Negeri Semarang, Semarang, Indonesia, Indonesia ,

(5) Department of Public Health, Universitas Negeri Semarang, Semarang, Indonesia, Indonesia

Abstract : 119

XML : 16 PDF : 168

XML

PDF

DOI : 10.20473/ijph.v21i1.2026.195-206

Login

Username *

Password *

[Forgot your password?](#)

☒ Keep me logged in

Login

[Register](#)

Accreditation Certificate

The Indonesian Journal of Public Health has been indexed by

Scopus®



Editorial Team



Prof. Kuntoro dr., MPH., Dr.PH

Editor in Chief

Department of Epidemiology, Biostatistics, Population Studies and Health Promotion,
Public Health Faculty, Universitas Airlangga, Indonesia

Interests: Biostatistics and Population

 0000-0002-0166-2394

kuntoro

7409844671



6061380



Dr. Lucia Yovita Hendrati, S.KM., M.Kes

Managing Editor

Department of Epidemiology, Biostatistics, Population Studies and Health Promotion,
Public Health Faculty, Universitas Airlangga, Indonesia

Interests: Epidemiology

 0000-0002-1287-0199

Lucia

57211793920



6054652



Dr. Nurul Fitriyah, S.KM., M.PH

Managing Editor

Department of Epidemiology, Biostatistics, Population Studies and Health Promotion,
Public Health Faculty, Universitas Airlangga, Indonesia

Interests: Maternal and Child Health, Reproductive Health

 0000-0001-6174-4394

Nurul Fitriyah

57212030024



6007948



Dr. Matthew Kelly

Editorial Board

Department of Applied Epidemiology, National Centre for Epidemiology and Population
Health, Australian National University, Australia

Interests: Population Health, Study Mortality

 0000-0001-7963-2139

Matthew Kelly

36125175000



-



Zhao Ni, Ph.D., BMed., RN., FAAN., FAHA

Editorial Board

Yale School of Nursing, Yale University, United States

Interests: Digital Health, Preventive Cardiology, HIV/AIDS Prevention, Nursing, Global
Health

 0000-0002-9185-9894

-

57190430263





Professor. Chung- Yi Li, Ph.D

Editorial Board

Department of Public Health, National Cheng Kung University, Taiwan

Interests: Environmental Epidemiology, Non-Communicable Disease Epidemiology, Life Course Epidemiology

 0000-0002-0321-8908

Chung-Yi Li

56829729900



Argita Dyah Salindri, S.KM., M.PH., Ph.D

Editorial Board

Stanford University School of Medicine, Stanford, United States of America

Interests: Epidemiology, Tuberculosis, Diabetes

 0000-0003-0962-0348

Argita Dyah Salindri

57193687955



Dr. Wan Ismahanisa Ismail, MLT, MSC Tran

Editorial Board

Department of Medical Laboratory Technology Faculty of Health Sciences Universiti Teknologi MARA (Pilau Pinang) Bertam Campus, Malaysia

Interests: Pharmacoepidemiology

 0000-0002-2711-0547

wan ismahanisa

57193530481



Associate Professor Dr. Muhammad Aziz Rahman, MBBS, MPH, CertGTC, GCHECTL, PhD

Editorial Board

Federation University Australia, Australia

Interests: Public Health, Tobacco, Health and Wellbeing, Disadvantaged Population, Non-Communicable Diseases

 0000-0003-1665-7966

Associate Professor Dr
Muhammad Aziz Rahman

55327627600



Associate Professor. Dr. Rajesh Ramasamy

Editorial Board

Department of Pathology, Faculty of Medicine & Health Science, Universiti Putra Malaysia, Malaysia

Interests: Mesenchymal Stem Cell, Immuno-modulation - Diabetes - Herbal Research

 0000-0003-4227-0458

Rajesh Ramasamy

26024442700





Associate Professor Dr Nor Afiah Mohd Zulkefli

Editorial Board

Department of Community Health, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia, Malaysia

Interests: Family Health



-

-

25724547000



-



Associate Professor. Dr. Normala Ibrahim MV BcH BAQ, MMed (Psych), PhD

Editorial Board

Departement of Psychiatry, Faculty of Medicine & Health Science, Universiti Putra Malaysia, Malaysia

Interests: Psychiatric Epidemiology



0000-0003-2868-9834

Normala Ibrahim

54414227000



-



Professor C. A. Kalpana, M.Sc., Bed., M.Phil., Ph.D

Editorial Board

Avinashilingam Institute for Home Science and Higher Education for Women, India

Interests: Obesity, Vitamin D, and Precision Nutrition, Digital Health Intervention, Machine Learning & Artificial Intelligence in Nutrition



-

Prof. C. A. Kalpana

-



-



Dr. Dian Kusuma, Scd., MPH

Editorial Board

City St George's, University of London, United Kingdom

Interests: Global Health, Health Systems, Health Policy, Health Economics, Non-Communicable Diseases



-

Dian Kusuma

57190047530



-



Yuniar Wardhani, S.KM., M.PH., Ph.D

Editorial Board

School of Public Health, Taipei Medical University, Taiwan, Province of China

Interests: Maternal and Child Health, Reproductive Health



-

Yuniar Wardani

-



-



Yashwant Pathak M.Pharm., EMBA, MS CM., Ph.D

Editorial Board

Faculty Affairs Collage of Pharmacy, University of South Florida, United States

Interests: Drug Delivery Systems and Nanotechnology



Yashwant Pathak

7004822104



Tito Yustiawan, MDD., MPH

Editorial Board

Departement of Health System, Faculty of Medical and Health Science, School of Population Health, University of Auckland, New Zealand

Interests: Health Administration and Policy, Hospital Administration, Hospital Management, Healthcare Management Information System



tito yustiawan

57210234979



6021142



Assoc. Prof. Dr. Supapron Chaigarun

Editorial Board

Ubon Ratchathani Rajabhat University, Thailand

Interests: Environmental Toxicology, Ecological Toxicology



-

50861236000



Dr. Agung Dwi Laksono, S.KM, M.Kes

Editorial Board

Center for Humanities, Health Policy, and Community Empowerment Research and Development Agency, Ministry of Health of the Republic of Indonesia, Indonesia

Interests: Healthcare Access, Health Disparities, Health Policy



0000-0002-9056-0399

Agung Dwi
Laksono

56429657100



6082273



Prof. Dr. Sri Sumarmi, SKM., M.Si

Editorial Board

Perhimpunan Sarjana Kesehatan Masyarakat (Society of Public Health Scholars), Indonesia

Interests: Nutrition, Public Health Nutrition, Nutrigenomic



0000-0002-7890-9578

Sri Sumarmi

55973957100



6004



Prof. Trias Mahmudiono, S.KM., M.PH (Nutr.), GCAS, P.hD

Editorial Board

World Public Health Nutrition Association, Indonesia

Interests: Nutrition, Public Health Nutrition, Nutrition Education, Nutritional Epidemiology, Double Burden

ID 0000-0002-3128-2173

Trias Mahmudiono

57189899256  5988526



Dr. Joni Haryanto, S.Kp., M.Si

Editorial Board

Faculty of Nursing, Airlangga University, Indonesia

Interests: Gerontology of Nursing

ID -

Joni Haryanto

57201187273

 5995607



Devi Aqilla Hastya Putri, S.KM

Administrative Assistant

Administration Staff of The Indonesian Journal of Public Health Universitas Airlangga, Indonesia

ID 0009-0007-0345-4162

-

-

 -



Farrah Febriastuti, S.Hum.

Administrative Assistant

Administration Staff Of The Indonesian Journal of Public Health Universitas Airlangga, Indonesia

ID 0009-0001-3019-813X

Farrah Febriastuti

-

 -

Login

Username *

Password *

[Forgot your password?](#)

☒ Keep me logged in

Login

Register

SENSITIVITY AND SPECIFICITY OF A NEW TOOL FOR EARLY DETECTION OF RISK FACTORS FOR NON COMMUNICABLE DISEASES IN URBAN WORKERS

Hildagardis Meliyani Erista Nai^{1*}, Riski Wulandari²

¹Department of Nutrition, Sekolah Tinggi Ilmu Kesehatan Panti Rapih Yogyakarta, Indonesia

²Department of Nursing, Sekolah Tinggi Ilmu Kesehatan Panti Rapih Yogyakarta, Indonesia

Corresponding author: Hildagardis Meliyani Erista Nai

Email: hildagardis_meliyani@stikespantirapih.ac.id

ABSTRACT

Introduction: Non communicable diseases (NCDs) have an impact on health, economy, and quality of life, and can reduce worker productivity. Approximately 41 million people die from NCDs every year, accounting for 74% of deaths worldwide. The Deteksi Dini Faktor Risiko Gizi dan Kesehatan (DDR-GizKes) is instrument designed for the early detection of nutritional and health risk factors related to NCDs that affect the productivity of urban workers. A screening test is essential part of this newly developed tool. **Aims:** This study aims to test the sensitivity and specificity of the DDR-GizKes instrument for detecting NCDs in urban workers. **Methods:** This study used a descriptive observational design with a cross-sectional approach. The population consisted of 277 teachers and staff in high schools in Yogyakarta who were selected using a cluster random sampling technique. Hypertension was used as the gold standard for the screening test. **Results:** The nutritional risk factor test had a sensitivity of 15.8% and a specificity of 94.2%. The positive predictive value (PPV) for the nutritional risk factor was 16.7% and the negative predictive value (NPV) was 93.8%. Meanwhile, the health risk factor test had a sensitivity of 15.8% and a specificity of 86.4%. The PPV for the health risk factor was 7.89% and the NPV was 93.3%. **Conclusion:** The DDR-GizKes instrument had low sensitivity but high specificity. Further research is necessary to establish the scoring system of the DDR-GizKes instrument in populations with a high prevalence of NCDs using a larger sample size.

Keywords: Nutrition, Health, DDR-GizKes, NCDs, Workers

INTRODUCTION

Noncommunicable diseases (NCDs) have an impact on the health, economy, and quality life of patients, families and communities (Rijal et al., 2019), reduce worker productivity (H et al., 2021), and place a burden on the healthcare system (World Health Organization, 2023b). Approximately 41 million people die from NCDs, accounting for 74% of deaths worldwide (World Health Organization, 2023b). The emergence of various NCDs, such as chronic obstructive pulmonary disease, cardiovascular disease, diabetes mellitus, and cancer, increases the burden of medical costs for patients and society (Kazibwe et al., 2021). Workers who are frequently sick or absent from work may impede the achievement of organizational goals, resulting in losses that also affect the national economy (World

Health Organization, 2019). NCDs are the leading cause of mortality and morbidity worldwide (Tolonen et al., 2023).

The prevalence of NCDs in Indonesia, including hypertension, stroke, cancer, chronic kidney disease, and diabetes mellitus, has increased from 2013 to 2018 (Ministry of Health, 2019). In the Special Region of Yogyakarta Province, the prevalence of NCDs such as diabetes mellitus and hypertension is also on the rise (Health Office of the Special Region of Yogyakarta, 2022). High blood pressure, also known as hypertension, is the most common risk factor for NCDs (Gupta and Xavier, 2018; Fuchs and Whelton, 2020). Hypertension affects approximately 1.28 billion people globally between the ages of 30 and 79 with the majority, or two-thirds, residing in low- and middle-income countries. One of the global goals for NCDs is to decrease the prevalence of

hypertension by 33% by 2030 (World Health Organization, 2023a).

In recent years, the prevalence of NCDs has increased due to rising economic status, changes in lifestyle, and the effects of modernization (Ismail and Sivadas, 2020). Risk factors of non-communicable diseases have also increased, including unhealthy eating patterns, lack of physical exercise, dangerous alcohol use, and exposure to tobacco smoke (Ministry of Health, 2019). The lifestyle of workers can also contribute to the development of NCDs, such as low physical activity and bad eating habits, including skipping breakfast, frequent snacking, consuming junk food, and eating few fruits and vegetables (Idris et al., 2021). Therefore, it is important to screen individuals at a productive age who are healthy or at risk of NCDs on a regular basis. In addition, early detection of risk factors is essential for disease prevention. Previous research has developed a risk self-assessment tool called *Deteksi Dini Faktor Risiko Gizi dan Kesehatan* (DDR-GizKes). The DDR-GizKes is an instrument in the form of a questionnaire designed for early detection of nutritional and health risk factors related to NCDs that affect the productivity of urban workers (Nai et al., 2023). A screening test is an essential part of this newly developed tool. Sensitivity and specificity analyses are commonly used to evaluate screening studies (Bujang and Adnan, 2016).

This study aims to address the problem of NCDs by evaluating the sensitivity and specificity of the DDR-GizKes instrument, which is designed for early detection of NCDs in urban workers. The key elements in the management of NCDs are detection, screening, and treatment (World Health Organization, 2023b). The DDR-GizKes instrument has been assessed for content validity by a panel of occupational nutrition experts and face validity by twenty employees with acceptable reliability. These validation did not include measuring sensitivity and

specificity (Nai et al., 2023). However, these validations did not include the evaluation of sensitivity and specificity of the DDR-GizKes instrument in comparison with the gold standard, which is crucial before applying this instrument to urban workers. In this study, hypertension was used as the gold standard for the DDR-GizKes instrument screening test. The incidence of hypertension is higher in individuals with cardiovascular disease and other serious illnesses (Gupta and Xavier, 2018). Therefore, detecting hypertension is critical to prevent the emergence of other NCDs. The novelty of this study is the assessment of the sensitivity and specificity of the DDR-GizKes instrument, which has not been previously evaluated (Nai et al., 2023). Therefore, this study aims to evaluate the sensitivity and specificity of the DDR-GizKes instrument for detecting NCDs in urban workers.

METHODS

Design, Location, and Time

This study used a descriptive observational design with a cross-sectional approach. This study was conducted in a number of senior high schools and vocational high schools in Yogyakarta City from July to October 2023.

Population, Sample Size, and Sampling Method

The population consisted of teachers and staff in a number of senior high schools and vocational high schools in Yogyakarta City who met the inclusion criteria. The participants were required to be between 20 and 40 years, willing to participate by signing informed consent, not currently on a certain diet, and not pregnant (for female participants). The sample size for the screening test was determined based on the Sample Size Calculation for Sensitivity and Specificity Analysis for Disease Prevalence of 5% to 20%. The estimated prevalence of hypertension in urban workers is 20%. The

null hypothesis (H_0) is 0.50 and the alternative hypothesis (H_a) is 0.60. The power is 0.804 and the p-value is 0.047. The minimum sample size required for positive disease is 50. In this study, the minimum sample size required was 249 participants (Bujang and Adnan, 2016). Ultimately, 277 participants who provided complete data were involved in this study.

The participants were selected using a cluster random sampling technique. The study was conducted in 72 senior high schools and vocational high schools in Yogyakarta City during the 2022/ 2023 semester (Ministry of Education, Culture, Research, and Technology of the Republic of Indonesia, 2023). The number of clusters was determined to be 14 schools, which is 20% of the total clusters (Nugraeni et al., 2023). Out of the 14 schools, the selected clusters were schools that gave permission to conduct this study and met the minimum sample size. Nine high schools selected as clusters, namely 1 State Senior High School, 2 State Senior High School, 7 State Senior High School, 11 State Senior High School, 2 State Vocational High School, 3 State Vocational High School, 4 State Vocational High School, Pangudi Luhur Senior High School, and Bintang Timur Catholic Senior High School. Teachers and staff in each school who met the inclusion criteria were involved in this study.

Variables

This study focuses on three main variables, namely sensitivity, specificity, and hypertension. Sensitivity is the proportion of positive test results among individuals who have the disease or the ability of the test to correctly identify those with the disease. Specificity is the proportion of negative test results among individuals who do not have the disease or the ability of the test to correctly identify those without the disease (Swift et al., 2020). A 2x2 table was used to calculate sensitivity and specificity.

		Status of person according to "gold standard"		
		Has the condition	Does not have the condition	
Result from screening test	Positive	a True positive	b False positive	Row entries for determining positive predictive value
	Negative	c False negative	d True negative	Row entries for determining negative predictive value
		Column entries for determining sensitivity	Column entries for determining specificity	

$$\text{Sensitivity} = [a / (a + c)] \times 100$$

$$\text{Specificity} = [d / (b + d)] \times 100$$

$$\text{Positive predictive value (PPV)} = [a / (a + b)] \times 100$$

$$\text{Negative predictive value (NPV)} = [d / (c + d)] \times 100.$$

Figure 1 .Calculation of Sensitivity, Specificity, PPV, and NPV (Trevethan, 2017)

Sensitivity and specificity tests were conducted on the DDR-GizKes instrument to evaluate nutritional and health risk factors. The DDR-GizKes instrument includes question items about risk factors for health and nutrition. There are 18 indicators for assessing NCDs in relation to nutritional risk factors and four indicators in relation to health risk factors. Each indicator is assigned a score of 0 to 10. A score of 10 indicates a category that is at risk for NCDs. The total score is the sum of all indicator scores for nutritional and health risk factors. The cut-off points for nutritional and health risk factors for detecting NCDs are 91 and 22, respectively. A nutritional risk factor score of 91 and higher and a health risk factor score of 22 and higher indicate a risk of developing NCDs (Nai et al., 2023).

The gold standard for the screening test in this study was hypertension with a blood pressure value 140/90 mmHg or greater (Ministry of Health, 2019). The blood pressure was measured using a sphygmomanometer.

The data collected from the participants included age range (20-30 years and 31-40 years), sex, education

level (elementary school dropout, elementary school, junior high school, senior high school, diploma, bachelor's degree, master's degree, and doctorate degree), income level (<IDR1,900,000, IDR 1,900,001 -3,800,000, IDR3,800,001-5,700,000, IDR5,700,001- 7,600,000, and >IDR7,600,000), and marital status (married, single, or divorced). The characteristics of the participants were collected using a structured questionnaire.

Data Collection

The data were collected by distributing the DDR-GizKes instrument and carrying out a physical examination. The characteristics of the participants were collected using a semi-structured questionnaire. A physical examination was carried out to collect data on blood pressure and question items in the DDR-GizKes instrument, including body weight, height, abdominal circumference, waist circumference, and hip circumference. The DDR-GizKes instrument was filled in directly by the participants with the assistance of an enumerator prior to taking the blood pressure measurement to prevent any influence from the results of the blood pressure measurement. This study involved 17 students as enumerators from the Nutrition Study Program and Nursing Study Program at Sekolah Tinggi Ilmu Kesehatan Panti Rapih Yogyakarta. The enumerators were trained in physical measurement and administration of the DDR-GizKes instrument.

Data Analysis

The data were analyzed using univariate analysis and screening tests. Univariate analysis was performed to describe the characteristics of the participants. Screening tests were carried out to analyze sensitivity, specificity, positive predictive value (PPV), and negative predictive value (NPV). STATA version 13 was used for the data analysis.

Ethical Approval

This study received approval from the Ethics Committee for Health Research of the Faculty of Health Science of Respati University of Yogyakarta with a certificate number 0145.3/FIKES/PL/VII/2023. Data collection in each school was carried out after obtaining permission from the school authorities. Informed consent was obtained from the participants prior to their participation in this study.

RESULTS

This study involved 277 participants. According to Table 1, the majority of the participants were aged between 31 and 40 years (57.4%), males (50.9%), and had a bachelor's degree (69.6%). In addition, the majority of the participants were married (69.3%) and had an income of between IDR1,900,001 and IDR3,800,000 (73.3%).

Table 1. Characteristics of the participants

Variable	Frequency	%
Age		
20- 30 years	118	42.6
31- 40 years	159	57.4
Sex		
Male	141	50.9
Female	136	49.1
Education Level		
Elementary school dropout	3	1.1
Elementary school	1	0.4

Variable	Frequency	%
Junior high school	1	0.4
Senior high school	41	14.8
Diploma	12	4.3
Bachelor's degree	193	69.6
Master's degree	26	9.4
Doctorate degree	0	0.0
Income Level		
< IDR 1,900,000	32	11.5
IDR 1,900,001- 3,800,000	203	73.3
IDR 3,800,001- 5,700,000	31	11.2
IDR 5,700,001- 7,600,000	8	2.9
>IDR7,600,000	3	1.1
Marital Status		
Married	192	69.3
Single/divorced	85	30.7
Total	277	100.0

Table 2. Sensitivity, specificity, PPV, and NPV of nutritional and health risk factors of the DDR-GizKes instrument

Parameters	Nutritional F actor	Health F actor
Prevalence : 6.9%		
True Positive	3	3
False Negative	16	16
True Negative	243	223
False Positive	15	35
Sensitivity	15.8%	15.8%
Specificity	94.2%	86.4%
Positive Likelihood Ratio	2.72	1.16
Negative Likelihood Ratio	0.89	0.97
Positive Predictive Value (PPV)	16.7%	7.89%
Negative Predictive Value (NPV)	93.8%	93.3%

Table 3. Comparison of area under the curve for nutrition and health risk factors

Variables	Area Under the Curve				p
	Area	95% Confidence interval		Standard Error	
		Lower Bound	Upper Bound		
Nutritional Factor	0.671	0.544	0.798	0.0647	0.004
Health Factor	0.402	0.263	0.540	0.071	

Table 2 shows that the prevalence of hypertension in the urban working population is 6.9%. The sensitivity test result for nutritional risk factors is 15.8%, indicating the ability of the nutritional risk factor test to yield positive results among urban workers who experienced hypertension is 15.8%. Meanwhile, while the remaining test results are negative (false negative). In comparison, the specificity test result for the nutritional risk factors is 94.2%, indicating that the ability of the nutritional risk factor test to yield negative results among urban workers who did not experience hypertension is 94.2%. Meanwhile, the remaining test results are positive (false positive). Furthermore, the positive predictive value (PPV) of the nutritional risk factor test for the prevalence of hypertension in urban workers is 16.7%, indicating that the true probability of experiencing hypertension if the nutritional risk factor test result is positive is 16.7%. In comparison, the negative predictive value (NPV) of the nutritional risk factor test for the prevalence of hypertension in urban workers is 93.8%, indicating that the true probability of not experiencing hypertension if the nutritional risk factor test result is negative is 93.8%.

Moreover, the sensitivity test result of the health risk factor test is 15.8%, indicating that the ability of the health risk factor test to yield positive results among urban workers with hypertension is 15.8%, while the remaining test results are negative (false negative). In contrast, the specificity test result of the health risk factor test is 86.4%, indicating that the ability of the health risk factor test to yield negative results among urban workers without hypertension is 86.4%, while the remaining test results are positive (false positive). The positive predictive value (PPV) of the health risk factor test for the prevalence of hypertension in urban workers is 7.89%, indicating that the true probability of experiencing hypertension if the health risk

factor test result is positive is 7.89%. In contrast, the negative predictive value (NPV) of the health risk factor test for the prevalence of hypertension in urban workers is 93.3%, indicating that the true probability of not experiencing hypertension if the health risk factor test result is negative is 93.3%.

Based on Table 3, Figure 2, and Figure 3, the area under the curve for the nutritional risk factor test is 0.671 (0.544 - 0.798) and for the health risk factor test was 0.402 (0.263 - 0.540) with a p-value of 0.004. Using a significance level (α) at 0.05, the ability of the nutritional and health risk factor tests to detect hypertension as the gold standard is different.




Figure 2.Area under the C curve of the Nutritional Risk F actor




Figure 3.Area under the Curve of the Health Risk Factor

DISCUSSION

This study aims to evaluate the *Deteksi Dini Faktor Risiko Gizi dan Kesehatan* (DDR-GizKes) instrument which has been developed and tested for content and face validity and reliability in previous research with good and acceptable results (Nai et al., 2023). In previous research, the DDR-GizKes instrument was developed based on theoretical insights regarding nutritional and health risk factors related to NCDs that can affect the productivity of urban workers. In this study, a scoring system was developed for each indicator of nutrition and health risk factors for early detection of NCDs in urban workers (Nai et al., 2023).

This study assessed the sensitivity and specificity of the DDR-GizKes instrument for early detection of NCDs in urban workers based on an established scoring system, specifically hypertension, using blood pressure measurement as the gold standard with a sphygmomanometer. The scoring system can be used to detect hypertension (Hatmalyakin et al., 2023). Previous research has developed a scoring system to predict the incidence of hypertension in the productive age group (Firmansyah et al., 2020). The DDR-GizKes instrument scoring system is differentiated for nutritional and health risk factors to detect NCDs. It includes 14 sub-variables and 18 indicators of nutritional risk factors, as well as six sub-variables and six indicators of health risk factors. To create the first DDR-GizKes instrument, 63 indicators were derived, 28 of which were related to nutritional risk factors and 35 to health risk factors. The total score for the nutritional risk factors was 145, which was the maximum score. Meanwhile, the total score for the health risk factors was 40, which was the total score of the four indicators (Nai et al., 2023).

The results of this study suggested that the sensitivity value for nutritional and health risk factors is low, while the specificity value is high. In addition, the

positive predictive value (PPV) for nutritional and health risk factors is low, while the negative predictive value (NPV) is high. A screening test is considered beneficial to public health if it can identify individuals at high risk of several NCDs with a sensitivity of at least 75% and a specificity of at least 65% (Gupta et al., 2017). Sensitivity and specificity indicate how well a test performs in comparison to the results from gold standards. On the other hand, the PPV and NPV indicate the effectiveness of a test in classifying individuals as having or not having the target condition based on the test findings (Trevethan, 2017).

The prevalence of a disease in a population can affect the sensitivity, specificity, PPV, and NPV of a screening tool (Hajian-Tilaki, 2014; Gogtay and Thatte, 2017; Trevethan, 2017) as well as the sample size (Bujang and Adnan, 2016). The prevalence of a disease varies from one population to another (Bujang and Adnan, 2016). A higher disease prevalence can result in higher sensitivity values and lower specificity values (Murad et al., 2023). In a population with a higher disease prevalence, the NPV is higher due to false positive results (Swift et al., 2020). Therefore, screening tests are carried out in populations with a higher risk of disease to reduce false positive results and increase the NPV (Bujang and Adnan, 2016).

This study examines the prevalence of hypertension in the urban working population. This study was conducted on urban workers due to their higher likelihood of nutritional and health risk factors related to NCDs (Idris et al., 2021). In addition, the age range for this study was between 20 and 40 years. Previous research shows that the incidence of hypertension is more prevalent among individuals aged between 50 and 60 years compared to younger individuals (Mularum et al., 2018). Moreover, age and blood pressure are positively correlated (Caraballo et al., 2021). Due to a number of anatomical and physiological modifications, the prevalence

of hypertension steadily increases with age (Jung et al., 2019). For individuals aged above 40 years, hypertension screening may be necessary if there is a growing trend in systolic blood pressure (Cheng et al., 2022). However, even tests with relatively high sensitivity and specificity may have low positive predictive value if the population prevalence is low. Therefore, to assess screening tests, it is also necessary to evaluate both the technical and population characteristics (Maxim et al., 2014). As a result, the DDR-GizKes instrument to be less sensitive in detecting NCDs in individuals aged under 50 years.

The prevalence of a disease affects the minimum sample size required for screening and diagnostic tests. Larger samples are necessary to detect higher levels of sensitivity for diseases with lower prevalence as well as higher levels of specificity for diseases with higher prevalence (Bujang and Adnan, 2016). Adequate sample size is the basis for determining the specificity and sensitivity of screening tests (Maxim et al., 2014). This study involved 277 participants for the DDR-GizKes instrument screening test, which is a relatively large sample. However, to achieve higher sensitivity and specificity, further research could include a larger sample size. The sensitivity and specificity of most screening tests can be determined with a minimum sample size of 300 individuals (Bujang and Adnan, 2016).

The DDR-GizKes instrument scoring system for predicting the risk of NCDs in urban workers also affects the level of sensitivity and specificity. The cut-off points for nutritional and health risk factors for NCD risk grouping may need to be reevaluated because sensitivity and specificity vary depending on the cut-off points. Specificity and sensitivity are negatively correlated, indicating that when sensitivity increases, specificity decreases, and vice versa. A threshold that prioritizes true negatives will result in more false negatives. Similarly, a threshold that prioritizes false positives will result in

more true positives (Gogtay and Thatte, 2017).

In addition, the scoring of several indicators of nutritional and health risk factors can be modified. For example, regularly skipping breakfast increases the risk of developing NCDs and is assigned a score of 10. Skipping breakfast has been associated with overweight and obesity (Wicherski et al., 2021), which in turn can impact the regulation or metabolism of the body, ultimately affecting weight management (Gibney et al., 2018). Meanwhile, obesity is a major contributor to serious NCDs (Zatońska et al., 2021). Moreover, insufficient sleep less than seven hours per night has been associated with an increased risk of NCDs (Mularum et al., 2018; Nurpratiwi et al., 2023). Therefore, this category is assigned a score of 10.

Changes in the scores for several indicators can alter the score range for nutritional and health risk factors. It is necessary to develop a scoring system to add the nutritional and health risk factor scores for the DDR-GizKes instrument. Therefore, further research is needed to develop the DDR-GizKes instrument scoring system for urban working populations with a more diverse age range. In addition, the gold standard should not only include the incidence of hypertension, but also the detection of NCDs with a high prevalence in urban workers, such as diabetes mellitus or coronary heart disease (Kuruvilla et al., 2023). For this reason, the research can use a case-control or prospective cohort design.

The results of the screening tests for the nutritional and health risk factors of the DDR-GizKes instrument indicated low sensitivity and high specificity. In conditions where funds and experts for diagnostic procedures are limited, screening tools with high specificity are necessary to reduce disease prevalence (Gupta et al., 2017). Therefore, screening tests are carried out on individuals who are most likely to be ill to minimize false positives. The DDR-GizKes instrument is a

screening tool for NCDs that is cheap, user-friendly, safe, and reliable. The indicators of the nutritional and health risk factors for the DDR-GizKes instrument to detect NCDs should be developed based on the balanced nutrition guidelines for the Indonesian society.

Limitations

One limitation of this study is that the population consisted of a young age group, namely individuals aged between 20 and 40 years, who have lower risk of hypertension compared to those above 40 years. In addition, the prevalence of hypertension in the urban working population is unknown, which may affect the sensitivity value. Furthermore, the NCDs studied only included hypertension. Although the sample size of this study met the required criteria, the sample size for the DDR-GizKes screening test still needs to be increased.

CONCLUSION

The DDR-GizKes instrument has been found to have low sensitivity and high specificity. Further investigation is required to evaluate its scoring system in urban worker populations with a high prevalence of NCDs using a larger sample size.

ACKNOWLEDGMENT

We would like to express our tremendous gratitude to the Directorate General of Higher Education, Research, and Technology of the Ministry of Education, Culture, Research, and Technology of the Republic of Indonesia for providing funding for this study (No. 0557/E5.5/AL.04/2023). We also extend our gratitude to all participants who took part in this study.

REFERENCES

Bujang, M. A., & Adnan, T. H. (2016).

Requirements for minimum sample size for sensitivity and specificity analysis. *Journal of Clinical and Diagnostic Research*, 10(10), YE01–YE06.

<https://doi.org/10.7860/JCDR/2016/18129.8744>

Caraballo, C., Wang, B., Angraal, S., Zhao, Y., Zheng, X., Lu, J., & Wu, C. (2021). Association of age and blood pressure among 3.3 million adults: Insights from China PEACE million persons project. *Journal of Hypertension*, 39(6), 1143–1154. <https://doi.org/10.1097/HJH.0000000000002793>

Cheng, W., Du, Y., Zhang, Q., Wang, X., He, C., He, J., Jing, F., Ren, H., Guo, M., Tian, J., & Xu, Z. (2022). Age-related changes in the risk of high blood pressure. *Frontiers in Cardiovascular Medicine*, 9(September), 1–10. <https://doi.org/10.3389/fcvm.2022.939103>

Firmansyah, Y., Ernawati, E., & Prawiro, E. L. (2020). Sistem Skoring Untuk Memprediksi Kejadian Hipertensi Pada Usia Produktif Di Kota Medan (Preliminary Study). *Jurnal Muara Sains, Teknologi, Kedokteran Dan Ilmu Kesehatan*, 4(1), 55. <https://doi.org/10.24912/jmstkik.v4i1.6013>

Fuchs, F. D., & Whelton, P. K. (2020). High Blood Pressure and Cardiovascular Disease. *Hypertension*, 75(2), 285–292. <https://doi.org/10.1161/HYPERTENSIONAHA.119.14240>

Gibney, M. J., Barr, S. I., Bellisle, F., Drewnowski, A., Fagt, S., Livingstone, B., Masset, G., Moreiras, G. V., Moreno, L. A., Smith, J., Vieux, F., Thielecke, F., & Hopkins, S. (2018). Breakfast in human nutrition: The international breakfast research initiative. *Nutrients*, 10(5), 1–12. <https://doi.org/10.3390/nu1005055>

- 9
- Gogtay, N. J., & Thatte, U. M. (2017). Statistical evaluation of diagnostic tests (part 1): Sensitivity, specificity, positive and negative predictive values. *Journal of Association of Physicians of India*, 65(JUNE), 80–84.
- Gupta, A., S, S., Aravindakshan, R., & Kekkar, R. (2017). A Pilot Retrospective Study Validating Noncommunicable Disease Risk Assessment Score (AMNRAS). *International Journal of Preventive Medicine*, 8. <https://doi.org/10.1016/j.ihj.2018.02.003>
- Gupta, R., & Xavier, D. (2018). Hypertension: The most important non communicable disease risk factor in India. *Indian Heart Journal*, 70(4), 565–572. <https://doi.org/10.1016/j.ihj.2018.02.003>
- H, M. S. S., Nisthar, S., & Nufile, A. A. M. (2021). *An analysis of the relationship between the productivity and the non-communicable diseases*. 14(3), 92–111.
- Hajian-Tilaki, K. (2014). Sample size estimation in diagnostic test studies of biomedical informatics. *Journal of Biomedical Informatics*, 48, 193–204. <https://doi.org/10.1016/j.jbi.2014.02.013>
- Health Service of Special Region of Yogyakarta. (2022). Health Profile of Special Region of Yogyakarta 2021. In *Health Service of Special Region of Yogyakarta*.
- Idris, I. B., Azit, N. A., Abdul Ghani, S. R., Syed Nor, S. F., & Mohammed Naw, A. (2021). A systematic review on noncommunicable diseases among working women. *Industrial Health*, 59(3), 146–160. <https://doi.org/10.2486/indhealth.2020-0204>
- Ismail, N. W., & Sivadas, S. (2020). Urban health and the prevalence of non-communicable diseases in Malaysia. *Malaysian Journal of Medicine and Health Sciences*, 16(2), 3–9.
- Jung, M. H., Yi, S. W., An, S. J., & Yi, J. J. (2019). Age-specific associations between systolic blood pressure and cardiovascular mortality. *Heart*, 105(14), 1070–1077. <https://doi.org/10.1136/heartjnl-2019-314697>
- Kazibwe, J., Tran, P. B., & Annerstedt, K. S. (2021). The household financial burden of non-communicable diseases in low- and middle-income countries: a systematic review. *Health Research Policy and Systems*, 19(1), 1–15. <https://doi.org/10.1186/s12961-021-00732-y>
- Kuruvilla, A., Mishra, S., & Ghosh, K. (2023). Prevalence and risk factors associated with non-communicable diseases among employees in a university setting: A cross-sectional study. *Clinical Epidemiology and Global Health*, 21(March), 101282. <https://doi.org/10.1016/j.cegh.2023.101282>
- Maxim, L. D., Niebo, R., & Utell, M. J. (2014). Screening tests: A review with examples. *Inhalation Toxicology*, 26(13), 811–828. <https://doi.org/10.3109/08958378.2014.955932>
- Ministry of Education Culture Research and Technology of the Republic of Indonesia. (2023). *Basic Education Data*. Ministry of Education Culture Research and Technology of the Republic of Indonesia. <https://dapo.kemdikbud.go.id/progress-sma/1/040000>
- Ministry of Health. (2019). *Baseline Health Research 2018 (Riskesdas 2018)*. Ministry of Health Republic of Indonesia.
- Mularum, S. N., Kusnanto, H., & Istiono, W. (2018). The Precision of

- Screening Questionnaires for Diabetes Mellitus Type 2 and Hypertension Compared with the Gold Standard in Primary Care. *Review of Primary Care Practice and Education (Kajian Praktik Dan Pendidikan Layanan Primer)*, 1(2), 75.
<https://doi.org/10.22146/rpcpe.36219>
- Murad, M. H., Lin, L., Chu, H., Hasan, B., Alsibai, R. A., Abbas, A. S., Mustafa, R. A., & Wang, Z. (2023). The association of sensitivity and specificity with disease prevalence: analysis of 6909 studies of diagnostic test accuracy. *CMAJ. Canadian Medical Association Journal*, 195(27), E925–E931.
<https://doi.org/10.1503/cmaj.221802>
- Nai, H. M. E., Estri, A. K., & Widiyanti, C. R. (2023). Developing a New Tool for Early Detection of the Nutritional and Health Risk Factors of Urban Workers' Productivity. *Kesmas*, 18(1), 41–49.
<https://doi.org/10.21109/kesmas.v18i1.6359>
- Nugraeni, T. A. E., Nai, H. M. E., & Maria, R. F. (2023). Hubungan Pola Konsumsi Makanan Cepat Saji dan Frekuensi Pemesanan Makanan Online dengan Obesitas Sentral pada Siswa SMA di Yogyakarta. *Amerta Nutrition*, 7(3), 413–420.
<https://doi.org/10.20473/amnt.v7i3.2023.413-420>
- Nurpratiwi, Hatmalyakin, D., Amaludin, M., Hidayat, U. R., Akbar, A., Arisandi, D., & Alfikrie, F. (2023). Early Detection of Hypertension (EDH) Skoring dalam Memprediksi Kejadian Hipertensi pada Usia Produktif. *Wiraraja Medika : Jurnal Kesehatan*, 13(1), 16–25.
<https://doi.org/10.24929/fik.v13i1.2503>
- Rijal, A., Adhikari, T. B., Khan, J. A. M., & Berg-Beckhoff, G. (2019). Correction: The economic impact of noncommunicable diseases among households in South Asia and their coping strategy: A systematic review (PLoS ONE (2018) 13:11 (e0205745)). DOI: 10.1371/journal.pone.0205745). *PLoS ONE*, 14(1), 1–23.
<https://doi.org/10.1371/journal.pone.0211588>
- Swift, A., Heale, R., & Twycross, A. (2020). What are sensitivity and specificity? *Evidence-Based Nursing*, 23(1), 2–5.
<https://doi.org/10.1136/ebnurs-2019-103225>
- Tolonen, H., Reinikainen, J., Zhou, Z., Härkänen, T., Männistö, S., Jousilahti, P., Paalanen, L., Lundqvist, A., & Laatikainen, T. (2023). Development of non-communicable disease risk factors in Finland: projections up to 2040. *Scandinavian Journal of Public Health*, 51(8), 1231–1238.
<https://doi.org/10.1177/14034948221110025>
- Trevethan, R. (2017). Sensitivity, Specificity, and Predictive Values: Foundations, Plausibilities, and Pitfalls in Research and Practice. *Frontiers in Public Health*, 5(November), 1–7.
<https://doi.org/10.3389/fpubh.2017.00307>
- Wicherski, J., Schlesinger, S., & Fischer, F. (2021). Observational Longitudinal Studies. *Nutrients*, 13(66492), 272.
<https://doi.org/10.3390/nu13010272>
- World Health Organization. (2019). Health, Decent Work and the Economy. *Policy Brief*, 18.
- World Health Organization. (2023a). *Hypertension*. World Health Organization.
- World Health Organization. (2023b). *Noncommunicable diseases*.
- Zatońska, K., Psikus, P., Basiak-Rasała, A., Stępnicka, Z., Gawęł-Dąbrowska,

D., Wołyniec, M., Gibka, J., Szuba, A., & Połtyn-Zaradna, K. (2021). Obesity and chosen non-communicable diseases in pure poland cohort study. *International*

Journal of Environmental Research and Public Health, 18(5), 1–10.
<https://doi.org/10.3390/ijerph18052701>



0.851852
Impact Factor


3028
Google Citations

Sinta 2
Current Accreditation

Google Scholar Garuda Website Editor URL

History Accreditation

Citation Per Year By Google Scholar



Journal By Google Scholar

	All	Since 2019
Citation	3028	2591



YAYASAN PANTI RAPIH
SEKOLAH TINGGI ILMU KESEHATAN PANTI RAPIH YOGYAKARTA

Jalan Tantular 401 Pringwulung, Condongcatur, Depok, Sleman, Yogyakarta
Telp.(0274) 518977, 542744 Fax (0274) 587143
Website : www.stikespantirapih.ac.id E-mail : stikespr@stikespantirapih.ac.id



SURAT TUGAS

Nomor: 373.K/STIKes-PR/A/III/2024

Ketua STIKes Panti Rapih Yogyakarta, dengan ini memberikan tugas kepada Saudara:

Nama: Ns. Riski Wulandari, M.Kep., Sp.Kep.M
NIDN: 0508039202
Status: Dosen Prodi Pendidikan Profesi Ners Program Profesi

Jurnal Publikasi	Jurnal
Sensitivity and Specificity of A New Tool For Early Detection of Risk Factors For Non Communicable Diseases in Urban Workers	The Indonesian Journal of Public Health (IJPH), Universitas Airlangga, Jurnal Nasional Terakreditasi Peringkat 2

Sehubungan dengan salah satu tugas dosen adalah melakukan publikasi ilmiah, maka dengan ini STIKes Panti Rapih Yogyakarta memberikan tugas pada Dosen tersebut untuk melakukan Publikasi Ilmiah pada Semester Genap Tahun Akademik 2023/2024.

Surat tugas ini berlaku mulai tanggal **1 Maret-31 Agustus 2024** dan untuk dipergunakan sebagaimana mestinya.

Yogyakarta, 1 Maret 2024

Ketua,

Yulia Wardani, MAN

SENSITIVITY AND SPECIFICITY OF A NEW TOOL FOR EARLY DETECTION OF RISK FACTORS FOR NONCOMMUNICABLE DISEASES IN URBAN WORKERS.pdf

by Logan Adams

Submission date: 05-Jul-2026 03:16PM (UTC+0900)

Submission ID: 2993302284

File name: SENSITIVITY_AND_SPECIFICITY_OF_A_NEW_TOOL_FOR_EARLY_DETECTION_OF_RISK_FACTORS_FOR_NONCOMMUNICABLE_DISEASES_IN_URBAN_WORKERS.pdf (443.24K)

Word count: 5704

Character count: 30616

**SENSITIVITY AND SPECIFICITY OF A NEW TOOL FOR EARLY
DETECTION OF RISK FACTORS FOR NON COMMUNICABLE DISEASES IN
URBAN WORKERS**1 **Hildagardis Meliyani Erista Nai^{1*}, Riski Wulandari²**¹Department of Nutrition, Sekolah Tinggi Ilmu Kesehatan Panti Rapih Yogyakarta, Indonesia²Department of Nursing, Sekolah Tinggi Ilmu Kesehatan Panti Rapih Yogyakarta, Indonesia

Corresponding author: Hildagardis Meliyani Erista Nai

Email: hildagardis_meliyani@stikespantirapih.ac.id

ABSTRACT

Introduction: Non communicable diseases (NCDs) have an impact on health, economy, and quality of life, and can reduce worker productivity. Approximately 41 million people die from NCDs every year, accounting for 74% of deaths worldwide. The Deteksi Dini Faktor Risiko Gizi dan Kesehatan (DDR-GizKes) is instrument designed for the early detection of nutritional and health risk factors related to NCDs that affect the productivity of urban workers. A screening test is essential part of this newly developed tool. **Aims:** This study aims to test the sensitivity and specificity of the DDR-GizKes instrument for detecting NCDs in urban workers. **Methods:** This study used a descriptive observational design with a cross-sectional approach. The population consisted of 277 teachers and staff in high schools in Yogyakarta who were selected using a cluster random sampling technique. Hypertension was used as the gold standard for the screening test. **Results:** The nutritional risk factor test had a sensitivity of 15.8% and a specificity of 94.2%. The positive predictive value (PPV) for the nutritional risk factor was 16.7% and the negative predictive value (NPV) was 93.8%. Meanwhile, the health risk factor test had a sensitivity of 15.8% and a specificity of 86.4%. The PPV for the health risk factor was 7.89% and the NPV was 93.3%. **Conclusion:** The DDR-GizKes instrument had low sensitivity but high specificity. Further research is necessary to establish the scoring system of the DDR-GizKes instrument in populations with a high prevalence of NCDs using a larger sample size.

Keywords: Nutrition, Health, DDR-GizKes, NCDs, Workers**INTRODUCTION**

1 Noncommunicable diseases (NCDs) have an impact on the health, economy, and quality life of patients, families and communities (Rijal et al., 2019), reduce worker productivity (H et al., 2021), and place a burden on the healthcare system (World Health Organization, 2023b). Approximately 41 million people die from NCDs, accounting for 74% of deaths worldwide (World Health Organization, 2023b). The emergence of various NCDs, such as chronic obstructive pulmonary disease, cardiovascular disease, diabetes mellitus, and cancer, increases the burden of medical costs for patients and society (Kazibwe et al., 2021). Workers who are frequently sick or absent from work may impede the achievement of organizational goals, resulting in losses that also affect the national economy (World

Health Organization, 2019). NCDs are the leading cause of mortality and morbidity worldwide (Tolonen et al., 2023).

The prevalence of NCDs in Indonesia, including hypertension, stroke, cancer, chronic kidney disease, and diabetes mellitus, has increased from 2013 to 2018 (Ministry of Health, 2019). In the Special Region of Yogyakarta Province, the prevalence of NCDs such as diabetes mellitus and hypertension is also on the rise (Health Office of the Special Region of Yogyakarta, 2022). High blood pressure, also known as hypertension, is the most common risk factor for NCDs (Gupta and Xavier, 2018; Fuchs and Whelton, 2020). Hypertension affects approximately 1.28 billion people globally between the ages of 30 and 79 with the majority, or two-thirds, residing 1 low- and middle-income countries. One of the global goals for NCDs is to decrease the prevalence of

1 Cite this as: Nai,H.M.E, and Wulandari, R, (2024). Sensitivity and Specificity of a New Tool for Early Detection of Risk Factors for Non-Communicable Diseases on Urban Workers.The Indonesian Journal of Public Health, 19(1), 94-105. <https://doi.org/10.20473/ijph.v19i1.2023.94-105>

1 hypertension by 33% by 2030 (World Health Organization, 2023a).

In recent years, the prevalence of NCDs has increased due to rising economic status, changes in lifestyle, and the effects of modernization (Ismail and Sivadas, 2020). Risk factors of non-communicable diseases have also increased, including unhealthy eating patterns, lack of physical exercise, dangerous alcohol use, and exposure to tobacco smoke (Ministry of Health, 2019). The lifestyle of workers can also contribute to the development of NCDs, such as low physical activity and bad eating habits, including skipping breakfast, frequent snacking, consuming junk food, and eating few fruits and vegetables (Idris et al., 2021). Therefore, it is important to screen individuals at a productive age who are healthy or at risk of NCDs on a regular basis. In addition, early detection of risk factors is essential for disease prevention. Previous research has developed a risk self-assessment tool called *Deteksi Dini Faktor Risiko Gizi dan Kesehatan* (DDR-GizKes). The DDR-GizKes is an instrument in the form of a questionnaire designed for early detection of nutritional and health risk factors related to NCDs that affect the productivity of urban workers (Nai et al., 2023). A screening test is an essential part of this newly developed tool. Sensitivity and specificity analyses are commonly used to evaluate screening studies (Bujang and Adnan, 2016).

This study aims to address the problem of NCDs by evaluating the sensitivity and specificity of the DDR-GizKes instrument, which is designed for early detection of NCDs in urban workers. The key elements in the management of NCDs are detection, screening, and treatment (World Health Organization, 2023b). The DDR-GizKes instrument has been assessed for content validity by a panel of occupational nutrition experts and face validity by twenty employees with acceptable reliability. These validation did not include measuring sensitivity and

specificity (Nai et al., 2023). However, these validations did not include the evaluation of sensitivity and specificity of the DDR-GizKes instrument in comparison with the gold standard, which is crucial before applying this instrument to urban workers. In this study, hypertension was used as the gold standard for the DDR-GizKes instrument screening test. The incidence of hypertension is higher in individuals with cardiovascular disease and other serious illnesses (Gupta and Xavier, 2018). Therefore, detecting hypertension is critical to prevent the emergence of other NCDs. The novelty of this study is the assessment of the sensitivity and specificity of the DDR-GizKes instrument, which has not been previously evaluated (Nai et al., 2023). Therefore, this study aims to evaluate the sensitivity and specificity of the DDR-GizKes instrument for detecting NCDs in urban workers.

METHODS

Design, Location, and Time

This study used a descriptive observational design with a cross-sectional approach. This study was conducted in a number of senior high schools and vocational high schools in Yogyakarta City from July to October 2023.

Population, Sample Size, and Sampling Method


The population consisted of teachers and staff in a number of senior high schools and vocational high schools in Yogyakarta City who met the inclusion criteria. The participants were required to be between 20 and 40 years, willing to participate by signing informed consent, not currently on a certain diet, and not pregnant (for female participants). The sample size for the screening test was **1** determined based on the Sample Size Calculation for Sensitivity and Specificity Analysis for Disease Prevalence of 5% to 20%. The estimated prevalence of hypertension in urban workers is 20%. The

null hypothesis (H0) is 0.50 and the alternative hypothesis (Ha) is 0.60. The power is 0.804 and the p-value is 0.047. The minimum sample size required for positive disease is 50. In this study, the minimum sample size required was 249 participants (Bujang and Adnan, 2016). Ultimately, 277 participants who provided complete data were involved in this study.

The participants were selected using a cluster random sampling technique. The study was conducted in 72 senior high schools and vocational high schools in Yogyakarta City during the 2022/ 2023 semester (Ministry of Education, Culture, Research, and Technology of the Republic of Indonesia, 2023). The number of clusters was determined to be 14 schools, which is 20% of the total clusters (Nugraeni et al., 2023). Out of the 14 schools, the selected clusters were schools that gave permission to conduct this study and met the minimum sample size. Nine high schools selected as clusters, namely 1 State Senior High School, 2 State Senior High School, 7 State Senior High School, 11 State Senior High School, 2 State Vocational High School, 3 State Vocational High School, 4 State Vocational High School, Pangudi Luhur Senior High School, and Bintang Timur Catholic Senior High School. Teachers and staff in each school who met the inclusion criteria were involved in this study.

Variables

This study focuses on three main variables, namely sensitivity, specificity, and hypertension. Sensitivity is the proportion of positive test results among individuals who have the disease or the ability of the test to correctly identify those with the disease. Specificity is the proportion of negative test results among individuals who do not have the disease or the ability of the test to correctly identify those without the disease (Swift et al., 2020). A 2x2 table was used to calculate sensitivity and specificity.



$$\text{Sensitivity} = [a / (a + c)] \times 100$$

$$\text{Specificity} = [d / (b + d)] \times 100$$

$$\text{Positive predictive value (PPV)} = [a / (a + b)] \times 100$$

$$\text{Negative predictive value (NPV)} = [d / (c + d)] \times 100.$$

Figure 1 .Calculation of Sensitivity, Specificity, PPV, and NPV (Trevethan, 2017)

Sensitivity and specificity tests were conducted on the DDR-GizKes instrument to evaluate nutritional and health risk factors. The DDR-GizKes instrument includes question items about risk factors for health and nutrition. There are 18 indicators for assessing NCDs in relation to nutritional risk factors and four indicators in relation to health risk factors. Each indicator is assigned a score of 0 to 10. A score of 10 indicates a category that is at risk for NCDs. The total score is the sum of all indicator scores for nutritional and health risk factors. The cut-off points for nutritional and health risk factors for detecting NCDs are 91 and 22, respectively. A nutritional risk factor score of 91 and higher and a health risk factor score of 22 and higher indicate a risk of developing NCDs (Nai et al., 2023).

The gold standard for the screening test in this study was hypertension with a blood pressure value 140/90 mmHg or greater (Ministry of Health, 2019). The blood pressure was measured using a sphygmomanometer.

The data collected from the participants included age range (20-30 years and 31-40 years), sex, education

level (elementary school dropout, elementary school, junior high school, senior high school, diploma, bachelor's degree, master's degree, and doctorate degree), income level (<IDR1,900,000, IDR 1,900,001 -3,800,000, IDR3,800,001-5,700,000, IDR5,700,001- 7,600,000, and >IDR7,600,000), and marital status (married, single, or divorced). The characteristics of the participants were collected using a structured questionnaire.

Data Collection

The data were collected by distributing the DDR-GizKes instrument and carrying out a physical examination. The characteristics of the participants were collected using a semi-structured questionnaire. A physical examination was carried out to collect data on blood pressure and question items in the DDR-GizKes instrument, including body weight, height, abdominal circumference, waist circumference, and hip circumference. The DDR-GizKes instrument was filled in directly by the participants with the assistance of an enumerator prior to taking the blood pressure measurement to prevent any influence from the results of the blood pressure measurement. This study involved 17 students as enumerators from the Nutrition Study Program and Nursing Study Program at Sekolah Tinggi Ilmu Kesehatan Panti Rapih Yogyakarta. The enumerators were trained in physical measurement and administration of the DDR-GizKes instrument.

1 Data Analysis

The data were analyzed using univariate analysis and screening tests. Univariate analysis was performed to describe the characteristics of the participants. Screening tests were carried out to analyze sensitivity, specificity, positive predictive value (PPV), and negative predictive value (NPV). STATA version 13 was used for the data analysis.

Ethical Approval

This study received approval from the Ethics Committee for Health Research of the Faculty of Health Science of Respati University of Yogyakarta with a certificate number 0145.3/FIKES/PL/VII/2023 . Data collection in each school was carried out after obtaining permission from the school authorities. Informed consent was obtained from the participants prior to their participation in this study.

RESULTS

This study involved 277 participants. According to Table 1, the majority of the participants were aged between 31 and 40 years (57.4%), males (50.9%), and had a bachelor's degree (69.6%). In addition, the majority of the participants were married (69.3%) and had an income of between IDR1,900,001 and IDR3,800,000 (73.3%).

Table 1. Characteristics of the participants

Variable	Frequency	%
Age		
20- 30 years	118	42.6
31- 40 years	159	57.4
Sex		
Male	141	50.9
Female	136	49.1
Education Level		
Elementary school dropout	3	1.1
Elementary school	1	0.4

Variable	Frequency	%
Junior high school	1	0.4
Senior high school	41	14.8
Diploma	12	4.3
Bachelor's degree	193	69.6
Master's degree	26	9.4
Doctorate degree	0	0.0
Income Level		
< IDR 1,900,000	32	11.5
IDR 1,900,001- 3,800,000	203	73.3
IDR 3,800,001- 5,700,000	31	11.2
IDR 5,700,001- 7,600,000	8	2.9
>IDR7,600,000	3	1.1
Marital Status		
Married	192	69.3
Single/divorced	85	30.7
Total	277	100.0

Table 2. Sensitivity, specificity, PPV, and NPV of nutritional and health risk factors of the DDR-GizKes instrument

Parameters	Nutritional F actor	Health F actor
Prevalence : 6.9%		
True Positive	3	3
False Negative	16	16
True Negative	243	223
False Positive	15	35
Sensitivity	15.8%	15.8%
Specificity	94.2%	86.4%
Positive Likelihood Ratio	2.72	1.16
Negative Likelihood Ratio	0.89	0.97
Positive Predictive Value (PPV)	16.7%	7.89%
Negative Predictive Value (NPV)	93.8%	93.3%

Table 3. Comparison of area under the curve for nutrition and health risk factors

Table 37. Comparison of area under the curve for nutritional and health risk factors					
Variables	Area Under the Curve				p
	Area	95% Confidence interval		Standard Error	
		Lower Bound	Upper Bound		
Nutritional Factor	0.671	0.544	0.798	0.0647	0.004
Health Factor	0.402	0.263	0.540	0.071	

1
Table 2 shows that the prevalence of hypertension in the urban working population is 6.9%. The sensitivity test result for nutritional risk factors is 15.8%, indicating the ability of the nutritional risk factor test to yield positive results among urban workers who experienced hypertension is 15.8%. Meanwhile, while the remaining test results are negative (false negative). In comparison, the specificity test result for the nutritional risk factors is 94.2%, indicating that the ability of the nutritional risk factor test to yield negative results among urban workers who did not experience hypertension is 94.2%. Meanwhile, the remaining test results are positive (false positive). Furthermore, the positive predictive value (PPV) of the nutritional risk factor test for the prevalence of hypertension in urban workers is 16.7%, indicating that the true probability of experiencing hypertension if the nutritional risk factor test result is positive is 16.7%. In comparison, the negative predictive value (NPV) of the nutritional risk factor test for the prevalence of hypertension in urban workers is 93.8%, indicating that the true probability of not experiencing hypertension if the nutritional risk factor test result is negative is 93.8%.

Moreover, the sensitivity test result of the health risk factor test is 15.8%, indicating that the ability of the health risk factor test to yield positive results among urban workers with hypertension is 15.8%, while the remaining test results are negative (false negative). In contrast, the specificity test result of the health risk factor test is 86.4%, indicating that the ability of the health risk factor test to yield negative results among urban workers without hypertension is 86.4%, while the remaining test results are positive (false positive). The positive predictive value (PPV) of the health risk factor test for the prevalence of hypertension in urban workers is 7.89%, indicating that the true probability of experiencing hypertension if the health risk

factor test result is positive is 7.89%. In contrast, the negative predictive value (NPV) of the health risk factor test for the prevalence of hypertension in urban workers is 93.3%, indicating that the true probability of not experiencing hypertension if the health risk factor test result is negative is 93.3%.

Based on Table 3, Figure 2, and Figure 3, the area under the curve for the nutritional risk factor test is 0.671 (0.544 - 0.798) and for the health risk factor test was 0.402 (0.263 - 0.540) with a p-value of 0.004. Using a significance level (α) at 0.05, the ability of the nutritional and health risk factor tests to detect hypertension as the gold standard is different.




Figure 2.Area under the C curve of the Nutritional Risk F factor




Figure 3.Area under the Curve of the Health Risk Factor

DISCUSSION

This study aims to evaluate the *Deteksi Dini Faktor Risiko Gizi dan Kesehatan* (DDR-GizKes) instrument which has been developed and tested for content and face validity and reliability in previous research with good and acceptable results (Nai et al., 2023). In previous research, the DDR-GizKes instrument was developed based on theoretical insights regarding nutritional and health risk factors related to NCDs that can affect the productivity of urban workers. In this study, a scoring system was developed for each indicator of nutrition and health risk factors for early detection of NCDs in urban workers (Nai et al., 2023).

This study assessed the sensitivity and specificity of the DDR-GizKes instrument for early detection of NCDs in urban workers based on an established scoring system, specifically hypertension, using blood pressure measurement as the gold standard with a sphygmomanometer. The scoring system can be used to detect hypertension (Hatmalyakin et al., 2023). Previous research has developed a scoring system to predict the incidence of hypertension in the productive age group (Firmansyah et al., 2020). The DDR-GizKes instrument scoring system is differentiated for nutritional and health risk factors to detect NCDs. It includes 14 sub-variables and 18 indicators of nutritional risk factors, as well as six sub-variables and six indicators of health risk factors. To create the first DDR-GizKes instrument, 63 indicators were derived, 28 of which were related to nutritional risk factors and 35 to health risk factors. The total score for the nutritional risk factors was 145, which was the maximum score. Meanwhile, the total score for the health risk factors was 40, which was the total score of the four indicators (Nai et al., 2023).

The results of this study suggested that the sensitivity value for nutritional and health risk factors is low, while the specificity value is high. In addition, the

positive predictive value (PPV) for nutritional and health risk factors is low, while the negative predictive value (NPV) is high. A screening test is considered beneficial to public health if it can identify individuals at high risk of several NCDs with a sensitivity of at least 75% and a specificity of at least 65% (Gupta et al., 2017). Sensitivity and specificity indicate how well a test performs in comparison to the results from gold standards. On the other hand, the PPV and NPV indicate the effectiveness of a test in classifying individuals as having or not having the target condition based on the test findings (Trevethan, 2017).

The prevalence of a disease in a population can affect the sensitivity, specificity, PPV, and NPV of a screening tool (Hajian-Tilaki, 2014; Gogtay and Thatte, 2017; Trevethan, 2017) as well as the sample size (Bujang and Adnan, 2016). The prevalence of a disease varies from one population to another (Bujang and Adnan, 2016). A higher disease prevalence can result in higher sensitivity values and lower specificity values (Murad et al., 2023). In a population with a higher disease prevalence, the NPV is higher due to false positive results (Swift et al., 2020). Therefore, screening tests are carried out in populations with a higher risk of disease to reduce false positive results and increase the NPV (Bujang and Adnan, 2016).

This study examines the prevalence of hypertension in the urban working population. This study was conducted on urban workers due to their higher likelihood of nutritional and health risk factors related to NCDs (Idris et al., 2021). In addition, the age range for this study was between 20 and 40 years. Previous research shows that the incidence of hypertension is more prevalent among individuals aged between 50 and 60 years compared to younger individuals (Mularum et al., 2018). Moreover, age and blood pressure are positively correlated (Caraballo et al., 2021). Due to a number of anatomical and physiological modifications, the prevalence

of hypertension steadily increases with age (Jung et al., 2019). For individuals aged above 40 years, hypertension screening may be necessary if there is a growing trend in systolic blood pressure (Cheng et al., 2022). However, even tests with relatively high sensitivity and specificity may have low positive predictive value if the population prevalence is low. Therefore, to assess screening tests, it is also necessary to evaluate both the technical and population characteristics (Maxim et al., 2014). As a result, the DDR-GizKes instrument to be less sensitive in detecting NCDs in individuals aged under 50 years.

The prevalence of a disease affects the minimum sample size required for screening and diagnostic tests. Larger samples are necessary to detect higher levels of sensitivity for diseases with lower prevalence as well as higher levels of specificity for diseases with higher prevalence (Bujang and Adnan, 2016). Adequate sample size is the basis for determining the specificity and sensitivity of screening tests (Maxim et al., 2014). This study involved 277 participants for the DDR-GizKes instrument screening test, which is a relatively large sample. However, to achieve higher sensitivity and specificity, further research could include a larger sample size. The sensitivity and specificity of most screening tests can be determined with a minimum sample size of 300 individuals (Bujang and Adnan, 2016).

The DDR-GizKes instrument scoring system for predicting the risk of NCDs in urban workers also affects the level of sensitivity and specificity. The cut-off points for nutritional and health risk factors for NCD risk grouping may need to be reevaluated because sensitivity and specificity vary depending on the cut-off points. Specificity and sensitivity are negatively correlated, indicating that when sensitivity increases, specificity decreases, and vice versa. A threshold that prioritizes true negatives will result in more false negatives. Similarly, a threshold that prioritizes false positives will result in

more true positives (Gogtay and Thatte, 2017).

In addition, the scoring of several indicators of nutritional and health risk factors can be modified. For example, regularly skipping breakfast increases the risk of developing NCDs and is assigned a score of 10. Skipping breakfast has been associated with overweight and obesity (Wicherski et al., 2021), which in turn can impact the regulation or metabolism of the body, ultimately affecting weight management (Gibney et al., 2018). Meanwhile, obesity is a major contributor to serious NCDs (Zatońska et al., 2021). Moreover, insufficient sleep less than seven hours per night has been associated with an increased risk of NCDs (Mularum et al., 2018; Nurpratiwi et al., 2023). Therefore, this category is assigned a score of 10.

Changes in the scores for several indicators can alter the score range for nutritional and health risk factors. It is necessary to develop a scoring system to add the nutritional and health risk factor scores for the DDR-GizKes instrument. Therefore, further research is needed to develop the DDR-GizKes instrument scoring system for urban working populations with a more diverse age range. In addition, the gold standard should not only include the incidence of hypertension, but also the detection of NCDs with a high prevalence in urban workers, such as diabetes mellitus or coronary heart disease (Kuruvilla et al., 2023). For this reason, the research can use a case-control or prospective cohort design.

The results of the screening tests for the nutritional and health risk factors of the DDR-GizKes instrument indicated low sensitivity and high specificity. In conditions where funds and experts for diagnostic procedures are limited, screening tools with high specificity are necessary to reduce disease prevalence (Gupta et al., 2017). Therefore, screening tests are carried out on individuals who are most likely to be ill to minimize false positives. The DDR-GizKes instrument is a

screening tool for NCDs that is cheap, user-friendly, safe, and reliable. The indicators of the nutritional and health risk factors for the DDR-GizKes instrument to detect NCDs should be developed based on the balanced nutrition guidelines for the Indonesian society.

Limitations

One limitation of this study is that the population consisted of a young age group, namely individuals aged between 20 and 40 years, who have lower risk of hypertension compared to those above 40 years. In addition, the prevalence of hypertension in the urban working population is unknown, which may affect the sensitivity value. Furthermore, the NCDs studied only included hypertension. Although the sample size of this study met the required criteria, the sample size for the DDR-GizKes screening test still needs to be increased.

CONCLUSION

The DDR-GizKes instrument has been found to have low sensitivity and high specificity. Further investigation is required to evaluate its scoring system in urban worker populations with a high prevalence of NCDs using a larger sample size.

ACKNOWLEDGMENT

We would like to express our tremendous gratitude to the Directorate General of Higher Education, Research, and Technology of the Ministry of Education, Culture, Research, and Technology of the Republic of Indonesia for providing funding for this study (No. 0557/E5.5/AL.04/2023). We also extend our gratitude to all participants who took part in this study.

REFERENCES

Bujang, M. A., & Adnan, T. H. (2016).

Requirements for minimum sample size for sensitivity and specificity analysis. *Journal of Clinical and Diagnostic Research*, 10(10), YE01–YE06.

<https://doi.org/10.7860/JCDR/2016/18129.8744>

Caraballo, C., Wang, B., Angraal, S., Zhao, Y., Zheng, X., Lu, J., & Wu, C. (2021). Association of age and blood pressure among 3.3 million adults: Insights from China PEACE million persons project. *Journal of Hypertension*, 39(6), 1143–1154. <https://doi.org/10.1097/HJH.0000000000002793>

Cheng, W., Du, Y., Zhang, Q., Wang, X., He, C., He, J., Jing, F., Ren, H., Guo, M., Tian, J., & Xu, Z. (2022). Age-related changes in the risk of high blood pressure. *Frontiers in Cardiovascular Medicine*, 9(September), 1–10. <https://doi.org/10.3389/fcvm.2022.939103>

Firmansyah, Y., Ernawati, E., & Prawiro, E. L. (2020). Sistem Skoring Untuk Memprediksi Kejadian Hipertensi Pada Usia Produktif Di Kota Medan (Preliminary Study). *Jurnal Muara Sains, Teknologi, Kedokteran Dan Ilmu Kesehatan*, 4(1), 55. <https://doi.org/10.24912/jmstkik.v4i1.6013>

Fuchs, F. D., & Whelton, P. K. (2020). High Blood Pressure and Cardiovascular Disease. *Hypertension*, 75(2), 285–292. <https://doi.org/10.1161/HYPERTENSIONAHA.119.14240>

Gibney, M. J., Barr, S. I., Bellisle, F., Drewnowski, A., Fagt, S., Livingstone, B., Masset, G., Moreiras, G. V., Moreno, L. A., Smith, J., Vieux, F., Thielecke, F., & Hopkins, S. (2018). Breakfast in human nutrition: The international breakfast research initiative. *Nutrients*, 10(5), 1–12. <https://doi.org/10.3390/nu1005055>

- 9
- Gogtay, N. J., & Thatte, U. M. (2017). Statistical evaluation of diagnostic tests (part 1): Sensitivity, specificity, positive and negative predictive values. *Journal of Association of Physicians of India*, 65(JUNE), 80–84.
- Gupta, A., S, S., Aravindakshan, R., & Kekkar, R. (2017). A Pilot Retrospective Study Validating Noncommunicable Disease Risk Assessment Score (AMNRAS). *International Journal of Preventive Medicine*, 8. <https://doi.org/10.1016/j.ihj.2018.02.003>
- Gupta, R., & Xavier, D. (2018). Hypertension: The most important non communicable disease risk factor in India. *Indian Heart Journal*, 70(4), 565–572. <https://doi.org/10.1016/j.ihj.2018.02.003>
- H, M. S. S., Nisthar, S., & Nufile, A. A. M. (2021). *An analysis of the relationship between the productivity and the non-communicable diseases*. 14(3), 92–111.
- Hajian-Tilaki, K. (2014). Sample size estimation in diagnostic test studies of biomedical informatics. *Journal of Biomedical Informatics*, 48, 193–204. <https://doi.org/10.1016/j.jbi.2014.02.013>
- Health Service of Special Region of Yogyakarta. (2022). Health Profile of Special Region of Yogyakarta 2021. In *Health Service of Special Region of Yogyakarta*.
- Idris, I. B., Azit, N. A., Abdul Ghani, S. R., Syed Nor, S. F., & Mohammed Nawi, A. (2021). A systematic review on noncommunicable diseases among working women. *Industrial Health*, 59(3), 146–160. <https://doi.org/10.2486/indhealth.2020-0204>
- Ismail, N. W., & Sivadas, S. (2020). Urban health and the prevalence of non-communicable diseases in Malaysia. *Malaysian Journal of Medicine and Health Sciences*, 16(2), 3–9.
- Jung, M. H., Yi, S. W., An, S. J., & Yi, J. J. (2019). Age-specific associations between systolic blood pressure and cardiovascular mortality. *Heart*, 105(14), 1070–1077. <https://doi.org/10.1136/heartjnl-2019-314697>
- Kazibwe, J., Tran, P. B., & Annerstedt, K. S. (2021). The household financial burden of non-communicable diseases in low- and middle-income countries: a systematic review. *Health Research Policy and Systems*, 19(1), 1–15. <https://doi.org/10.1186/s12961-021-00732-y>
- Kuruvilla, A., Mishra, S., & Ghosh, K. (2023). Prevalence and risk factors associated with non-communicable diseases among employees in a university setting: A cross-sectional study. *Clinical Epidemiology and Global Health*, 21(March), 101282. <https://doi.org/10.1016/j.cegh.2023.101282>
- Maxim, L. D., Niebo, R., & Utell, M. J. (2014). Screening tests: A review with examples. *Inhalation Toxicology*, 26(13), 811–828. <https://doi.org/10.3109/08958378.2014.955932>
- Ministry of Education Culture Research and Technology of the Republic of Indonesia. (2023). *Basic Education Data*. Ministry of Education Culture Research and Technology of the Republic of Indonesia. <https://dapo.kemdikbud.go.id/progress-sma/1/040000>
- Ministry of Health. (2019). *Baseline Health Research 2018 (Riskesdas 2018)*. Ministry of Health Republic of Indonesia.
- Mularum, S. N., Kusnanto, H., & Istiono, W. (2018). The Precision of

- Screening Questionnaires for Diabetes Mellitus Type 2 and Hypertension Compared with the Gold Standard in Primary Care. *Review of Primary Care Practice and Education (Kajian Praktik Dan Pendidikan Layanan Primer)*, 1(2), 75.
<https://doi.org/10.22146/rpcpe.36219>
- Murad, M. H., Lin, L., Chu, H., Hasan, B., Alsibai, R. A., Abbas, A. S., Mustafa, R. A., & Wang, Z. (2023). The association of sensitivity and specificity with disease prevalence: analysis of 6909 studies of diagnostic test accuracy. *CMAJ. Canadian Medical Association Journal*, 195(27), E925–E931.
<https://doi.org/10.1503/cmaj.221802>
- Nai, H. M. E., Estri, A. K., & Widiyanti, C. R. (2023). Developing a New Tool for Early Detection of the Nutritional and Health Risk Factors of Urban Workers' Productivity. *Kesmas*, 18(1), 41–49.
<https://doi.org/10.21109/kesmas.v18i1.6359>
- Nugraeni, T. A. E., Nai, H. M. E., & Maria, R. F. (2023). Hubungan Pola Konsumsi Makanan Cepat Saji dan Frekuensi Pemesanan Makanan Online dengan Obesitas Sentral pada Siswa SMA di Yogyakarta. *Amerta Nutrition*, 7(3), 413–420.
<https://doi.org/10.20473/amnt.v7i3.2023.413-420>
- Nurpratiwi, Hatmalyakin, D., Amaludin, M., Hidayat, U. R., Akbar, A., Arisandi, D., & Alfikrie, F. (2023). Early Detection of Hypertension (EDH) Skoring dalam Memprediksi Kejadian Hipertensi pada Usia Produktif. *Wiraraja Medika : Jurnal Kesehatan*, 13(1), 16–25.
<https://doi.org/10.24929/fik.v13i1.2503>
- Rijal, A., Adhikari, T. B., Khan, J. A. M., & Berg-Beckhoff, G. (2019). Correction: The economic impact of noncommunicable diseases among households in South Asia and their coping strategy: A systematic review (PLoS ONE (2018) 13:11 (e0205745)). DOI: 10.1371/journal.pone.0205745).
PLoS ONE, 14(1), 1–23.
<https://doi.org/10.1371/journal.pone.0211588>
- Swift, A., Heale, R., & Twycross, A. (2020). What are sensitivity and specificity? *Evidence-Based Nursing*, 23(1), 2–5.
<https://doi.org/10.1136/ebnurs-2019-103225>
- Tolonen, H., Reinikainen, J., Zhou, Z., Härkänen, T., Männistö, S., Jousilahti, P., Paalanen, L., Lundqvist, A., & Laatikainen, T. (2023). Development of non-communicable disease risk factors in Finland: projections up to 2040. *Scandinavian Journal of Public Health*, 51(8), 1231–1238.
<https://doi.org/10.1177/14034948221110025>
- Trevethan, R. (2017). Sensitivity, Specificity, and Predictive Values: Foundations, Plausibilities, and Pitfalls in Research and Practice. *Frontiers in Public Health*, 5(November), 1–7.
<https://doi.org/10.3389/fpubh.2017.00307>
- Wicherski, J., Schlesinger, S., & Fischer, F. (2021). Observational Longitudinal Studies. *Nutrients*, 13(66492), 272.
<https://doi.org/10.3390/nu13010272>
- World Health Organization. (2019). Health, Decent Work and the Economy. *Policy Brief*, 18.
- World Health Organization. (2023a). *Hypertension*. World Health Organization.
- World Health Organization. (2023b). *Noncommunicable diseases*.
- Zatońska, K., Psikus, P., Basiak-Rasała, A., Stępnicka, Z., Gawel-Dabrowska,

D., Wołynec, M., Gibka, J., Szuba, A., & Połtyn-Zaradna, K. (2021). Obesity and chosen non-communicable diseases in pure poland cohort study. *International*

Journal of Environmental Research and Public Health, 18(5), 1–10.
<https://doi.org/10.3390/ijerph18052701>

SENSITIVITY AND SPECIFICITY OF A NEW TOOL FOR EARLY DETECTION OF RISK FACTORS FOR NONCOMMUNICABLE DISEASESIN URBAN WORKERS.pdf

ORIGINALITY REPORT

78%
SIMILARITY INDEX

78%
INTERNET SOURCES

87%
PUBLICATIONS

53%
STUDENT PAPERS

PRIMARY SOURCES

1 repository.unar.ac.id
Internet Source

Exclude quotes
Exclude bibliography

On
Off

Exclude matches

< 10%